


Mixing instructions for  
**VARIBAR® THIN LIQUID**  
(barium sulfate) for oral suspension

The VARIBAR THIN LIQUID powder must be reconstituted prior to administration by a healthcare provider according to the following instructions:



(Item # 900001)




**STEP 1:** Add water to the fill line on the bottle label. Replace the lid securely.



**STEP 2:** Invert the bottle and tap with fingers to mix the powder into the water.



**STEP 3:** Shake vigorously for 30 seconds.



**STEP 4:** Let stand for 5 minutes.



**STEP 5:** Re-fill with water to the Fill Line on the bottle label and re-shake thoroughly.



**STEP 6:** Once reconstituted, write the “discard after” date on the immediate container label.

- After reconstitution, store in refrigerator at 2°C to 8°C (36°F to 46°F) for up to 72 hours.
- Reconstitution yields approximately 300 mL of VARIBAR THIN LIQUID oral suspension containing 0.4 grams of barium sulfate per mL (40% w/v) and should be homogeneous and white to lightly colored.

For dosing and other prescribing information see prescribing information on the following pages








# Varibar® (barium sulfate)

Mixing instructions for  
**VARIBAR® THIN LIQUID**  
(barium sulfate)  
for oral suspension



(Item #900001)

VARIBAR is manufactured by E-Z-EM Canada Inc., for E-Z-EM, Inc., a subsidiary of Bracco Diagnostics Inc., Princeton, NJ 08540. VARIBAR is a registered trademark of E-Z-EM, Inc.

Bracco Diagnostics Inc.  
510 Carnegie Center | Suite 300  
Princeton, NJ 08540 USA

Phone: 609-514-2200  
Toll-Free: 1-800-631-5245 (U.S. only)  
Fax: 609-514-2424

©2025 Bracco Diagnostics Inc. All Rights Reserved. US-VARHL-2500001 01/25

**UNLOCKING THE  
INVISIBLE**

